Mathematics for Machine Learning (2025 Fall) Assignment 02*

Chuang-Chieh Lin

Department of Computer Science and Engineering, National Taiwan Ocean University

1. (20%) Diagonalize

$$\begin{bmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{bmatrix} = \mathbf{P} \mathbf{D} \mathbf{P}^{\top}$$

such that ${f P}$ consists of orthonormal column vectors.

- 2. (20%) Suppose that $\mathbf{y} = (y_1, y_2 \dots, y_n) \in \mathbb{R}^n$ is the observation and $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ is a prediction from a model such that $\sum_{i=1}^n x_i = \sum_{j=1}^n y_j = 1$. What is $\frac{\partial}{\partial \mathbf{x}} D_{\mathrm{KL}}(\mathbf{x} || \mathbf{y})$?
- 3. (20%) Given $\mathbf{x} \in \mathbb{R}^n$, please compute $\frac{d}{d\mathbf{x}}(2\mathbf{x})$.
- 4. (20%) Compute the derivative f'(x) of the (univariate) logistic sigmoid

$$f(x) = \frac{1}{1 + \exp(-x)}.$$

5. (20%) Given the formula

$$\frac{\partial \mathbf{x}^{\top} \mathbf{B} \mathbf{x}}{\partial \mathbf{x}} = \mathbf{x}^{\top} (\mathbf{B} + \mathbf{B}^{\top})$$

for a square matrix **B**, compute the gradient

$$\frac{\partial}{\partial \mathbf{s}} \big((\mathbf{x} - \mathbf{A}\mathbf{s})^{\top} \mathbf{A} \mathbf{A}^{\top} (\mathbf{x} - \mathbf{A}\mathbf{s}) + \|\mathbf{s}\|^2 \big)$$

where $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{s} \in \mathbf{R}^{n \times 1}$ and $\mathbf{x} \in \mathbf{R}^{m \times 1}$.

^{*} List the required intermediate steps next to each problem. Note that any answers generated directly by AI are invalid for this assignment.